

References

E-17

1. Antonini A, Funch VSC, Boyd JT, et al. Effect of Levodopa-Carbidopa Intestinal Gel on Dyskinesia in Advanced Parkinson's Disease Patients. *Move Disord*. 2016;530-536.
2. Noridian Healthcare Solutions, LLC. L33794. Effective 10/01/2015. Revised 01/01/2017.
3. Hayes, Inc. Hayes clinical research response. *Patient-Controlled Analgesia Infusion Pumps*. Lansdale,Pa; Hayes, Inc; Published 04/08/2019.
4. CMS Manual System, Pub 100-3 Medical National coverage Determination Manual, Chapter 1, Section 280.14. Infusion Pumps.
5. Hobbs JG, Ryan KR, Ritchie B, Sluggett, et al. Protocol for a randomized crossover trial to evaluate patient and nurse satisfaction with electronic and elastomeric portable infusion pumps for the continuous administration of antibiotic therapy in the home: The comparing home infusion devices (CHID) study. *BMJ* 2017;7e016763.
6. Perks SJ, Pain T, Franklin R. Total intended antibiotic delivery related to drug concentration affecting the flow rate of elastomeric devices used in outpatient parenteral therapy (OPAT). *J Pharm Prac and Research*. 2019; 49: 349-355.