

## References

### E-34

1. Cui L, Liu H, Sun L. Multidisciplinary respiratory rehabilitation in combination with non-invasive positive pressure ventilation in the treatment of elderly patients with severe chronic obstructive pulmonary disease. *Pak J Med Sci.* 2019; 35(2).
2. Nasralla MLS, Bolzan DW, Lage YG, et al. Extended-time of noninvasive positive Pressure ventilation improves tissue perfusion after coronary artery bypass surgery: A randomized clinical trial. *Braz J Cardiovasc Surg.* 2018; 33(3):250-257.
3. Ishfaq N, Gul N, Zaka N. Outcome of early use of non-invasive positive pressure ventilation in patients with acute exacerbation of chronic obstructive pulmonary disease. *Pak J Med Sci.* 2019;35(6):1488-1492.
4. Dorst J, Behrendt G, Ludolph. Noninvasive ventilation and hypercapnia-associated symptoms I amyotrophic lateral sclerosis. *Acta Neurol Scand.* 2019;139:128–134.
5. Heiman-Patterson T, Cudkowicz M, De Carvalho M, et al. Understanding the use of HIV in ALS: Results of an international ALS specialist survey. *ALS and Frontotemporal Deg* 2018;19:331–341.
6. Benditt J. Respiratory care of patients with neuromuscular disease. *Respir Care.* 2019;64(6):679–688.