

References

E-45

1. Eftekharsadat B, Babaei-Ghazani A, Habibzadeh A, et al. Efficacy of action potential simulation and interferential therapy in the rehabilitation of patients with knee osteoarthritis. *Ther Adv Musculoskelet Dis*. 2015;7(3):67-75.
2. Vitton V, Benezech A, Honore S, et al. CON-COUR study: Interferential therapy in the treatment of chronic constipation in adults: Study protocol for a randomized controlled trial. 2015;16:234.
3. Qaseem A, Wilt T, McLean R, Forciea M. Noninvasive treatments for acute, subacute, and chronic low back pain: A clinical practice guideline from the American College of Physicians. *Ann Intern Med*. 2017;166(7):514-530.
4. Zeng C, Li H, Yang T, et al. Electrical stimulation for pain relief in knee osteoarthritis: Systematic review and network meta-analysis. *Osteoarthritis Cartilage*. 2015;23(2):189-202.
5. Moore, JS, Gibson, PR, Burgell, RE. Neuromodulation via interferential electrical stimulation as a novel therapy in gastrointestinal motility disorders. *J Neurogastroenterol Motil*. 2018;30;24(1):19-29.
6. National Institute for Health and Care Excellence: Clinical Guidelines (NICE), 2016. Low back pain and sciatica in over 16s: Assessment and management. Accessed on June 11, 2018.
7. Lindblad K, Bergkvist L, Johansson A-C. Evaluation of the treatment of chronic chemotherapy-induced peripheral neuropathy using long-wave diathermy and interferential currents: A randomized controlled trial. *Supportive Care in Cancer*. 2016;24(6):2523-2531.
8. Mohite SA, Shinde SB. Effect of early intervention with spinal isometric exercises in acute lumbar intervertebral disc prolapse. *Indian J Physiot Occup Ther*. 2019;13(2):170175.
9. Corrêa JB, Costa LOP, Oliveira NTB, et al. Effects of the carrier frequency of interferential current on pain modulation and central hypersensitivity in people with chronic nonspecific low back pain: A randomized placebo-controlled trial. *European J Of Pain*. 2016;20(10):1653-1666.

10. Mahmoud MM, El-Nahas NG, Hamed MH, Louis NN. Interferential electrical stimulation versus pulsed electro-magnetic field in management of intermittent claudication. *Egy J Hosp Med.* 2020;80: 654-661.
11. Singh SK, Agrawal R, Akbani R. Comparison of the effect of high voltage pulsed current versus interferential therapy on pain and womac in patients with knee osteoarthritis. *Ind J of Physio Occup Ther.* 2019;13(4).