

Reference

M-23

1. Hall, M., Wenner, J., Oberg, S. The postprandial acid pocket does not extend into most distal esophagus: Evidence from pH studies performed immediately above the squamocolumnar junction. *Scandinavia Journal of Gastroenterology*. 2014;49(1):15-22.
2. Kleiman, D., Beninato, T., Bosworth, BP., et al. Early referral for esophageal pH monitoring is more cost-effective than prolonged empiric trials of proton-pump inhibitors for suspected gastroesophageal reflux disease. *Journal of Gastrointestinal Surgery*. 2014;18(1):26-33.
3. Fitzgerald R, di Pietro M, Raquinath K, et al. British Society of Gastroenterology guidelines on the diagnosis and management of Barrett's esophagus. *Gut*. 2014;63(1):7-42.
4. Richter J, Pandolfino J, Vela M, et al. Utilization of wireless pH monitoring technologies: A summary of the proceedings from the Esophageal Diagnostic Working Group. *Disease of the Esophagus*. 2013;26:755-765.
5. Kleiman D, Sporn M, Beninato T, Metz Y, et al. Early referral for 24-h esophageal pH monitoring may prevent unnecessary treatment with acid-reducing medications. *Surgical Endoscopy*. 2013; 27:1302-1309.
6. Kavitt RT, Yuksel E, Slaughter J, et al. The role of impedance monitoring in patients with extraesophageal symptoms. *The Laryngoscope*. 2013;123(10):2463-2468.
7. Katz P, Gerson L, Vela M. Guidelines for the diagnosis and management of gastroesophageal reflux disease. *American Journal of Gastroenterology*. 2013;108:308 – 328.
8. Yang X, Gan T, Wang L, et al. Wireless esophageal pH capsule for patients with gastroesophageal reflux disease: A multicenter clinical study. *World Journal of Gastroenterology*. 2014; 20(40):14865-14874.
9. Shin M. Esophageal pH and combined impedance-pH monitoring in children. *Pediatric Gastroenterology, Hepatology & Nutrition*. 2014;17(1):13-22.
10. Iluyomade A, Olowoyeye A, Fadahunsi O, et al. Interference with daily activities and major adverse events during esophageal pH monitoring with bravo wireless capsule versus conventional intranasal catheter: A systematic review of randomized controlled trials. *Disease of the Esophagus*. 2017;30(3): 1-9.
11. Cresi F, Locatelli E, Marinaccio C, et al. Prognostic values of multichannel intraluminal impedance and pH monitoring in newborns with symptoms of gastroesophageal reflux disease. *The Journal of Pediatrics*. 2013;162(4):770-5
12. Katz P, Gerson L, Vela M. Guidelines for the diagnosis and management of gastroesophageal reflux disease. *American Journal of Gastroenterology*. 2013; 108(3):308-28.
13. Salvatore S, Salvatoni A, Van Berkel M, et al. Esophageal impedance baseline is age dependent. *J Pediatr Gastroenterol Nutr*. 2013;57(4):506-513.
14. National Institute for Health and Care Excellence (NICE). Gastro-oesophageal reflux disease in children and young people: diagnosis and management. 2015. Accessed August 28, 2018.
15. Singendonk MMJ, Benninga MA, van Wijk MP. Reflux monitoring in children. *Neurogastroenterology And Motility: The Official Journal Of The European Gastrointestinal Motility Society*. 2016;28(10):1452- 1459.
16. Kahrilas P, Shaheen N, Vaezi M. American Gastroenterological Association medical Position Statement on the Management of Gastroesophageal Reflux Disease. 2018;135(4):1383-1391.

17. Iluyomade A, Olowoyeye A, Fadahunsi O, et al. Interference with daily activities and major adverse events during esophageal pH monitoring with bravo wireless capsule versus conventional intranasal catheter: a systematic review of randomized controlled trials. *Diseases Of The Esophagus: Official Journal Of The International Society For Diseases Of The Esophagus*. 2017;30(3):1-9.