

References

S-157

1. American Orthopaedic Foot & Ankle Society. Extracorporeal Shock Wave Therapy. 2017.
2. Kertzman P, Császár N, Furia J, Schmitz C. Radial extracorporeal shock wave therapy is efficient and safe in the treatment of fracture non unions of superficial bones: A retrospective case series. *J Orthop Surg Res.* 2017;12(1):164
3. Malliaropoulos N, Thompson D, Lohrer H, et al. Individualised radial extracorporeal shock wave therapy (rESWT) for symptomatic calcific shoulder tendinopathy: A retrospective clinical study. *BMC Musculoskeletal Disord.* 2017;18(1):513.
4. Liao C-D, Xie G-M, Tsao J-Y, et.al. Efficacy of extracorporeal shock wave therapy for knee tendinopathies and other soft tissue disorders: A meta-analysis of randomized controlled trials. *BMC Musculoskeletal Disord.* 2018;19(1):278.
5. Ibrahim MI, Donatelli RA, Hellman M, Hussein AZ, Furia JP, Schmitz C. Long-term results of radial extracorporeal shock wave treatment for chronic plantar fasciopathy: A prospective, randomized, placebo-controlled trial with two years follow- *J Orthop Res.* 2017;35(7):1532-8.
6. Lai TW, Ma HL, Lee MS, Chen PM, Ku MC. Ultrasonography and clinical outcome comparison of extracorporeal shock wave therapy and corticosteroid injections for chronic plantar fasciitis: a randomized controlled trial. *J Musculoskeletal Neuronal Interact.* 2018;18(1):47.
7. Lizis P. Analgesic effect of extracorporeal shock wave therapy versus ultrasound therapy in chronic tennis elbow. *J Phys Ther Sci.* 2015;27(8):2563-2567.
8. Alessio-Mazzola M, Repetto I, Biti B, et al. Autologous US-guided PRP injection versus US-guided focal extracorporeal shock wave therapy for chronic lateral epicondylitis: a minimum of 2-year follow-up retrospective comparative study. *J Orthop Surg.* 2018;26(1):2309499017749986.
9. Li S, Wang K, Sun H, et al. Clinical effects of extracorporeal shock-wave therapy and ultrasound-guided local corticosteroid injections for plantar fasciitis in adults: A meta-analysis of randomized controlled trials. *Medicine.* 2018;97(50).
10. Williams HL, Jones SA, Lyons C, Wilson C, Ghandour A. Refractory patella tendinopathy with failed conservative treatment—shock wave or arthroscopy? *J Orthop Surg.* 2017;25(1):2309499016684700.
11. Liao CD, Xie GM, Tsao JY, Chen HC, Liou TH. Efficacy of extracorporeal shock wave therapy for knee tendinopathies and other soft tissue disorders: a

- meta-analysis of randomized controlled trials. *BMC Musculoskeletal Disord.* 2018;19(1):278.
12. Zhang Q, Liu L, Sun W, Gao F, Cheng L, Li Z. Extracorporeal shockwave therapy in osteonecrosis of femoral head: a systematic review of now available clinical evidences. *Medicine.* 2017;96(4).
 13. Furia JP, Rompe JD, Maffulli N, Cacchio A, Schmitz C. Radial extracorporeal shock wave therapy is effective and safe in chronic distal biceps tendinopathy. *Clin J Sport Med.* 2017;27(5):430-7.
 14. Hao Y, Guo H, Xu Z, et al. Meta-analysis of the potential role of extracorporeal shockwave therapy in osteonecrosis of the femoral head. *J Orthop Surg Res.* 2018;13(1):166.
 15. Ko JY, Siu KK, Wang FS, et al. The therapeutic effects of extracorporeal shock wave therapy (ESWT) on the rotator cuff lesions with shoulder stiffness: A prospective randomized study. *BioMed Res Int.* 2020;2020.
 16. Malliaropoulos N, Thompson D, Meke M, et al. Individualised radial extracorporeal shock wave therapy (rESWT) for symptomatic calcific shoulder tendinopathy: a retrospective clinical study. *BMC Musculoskeletal Disord.* 2017;18(1):1-7.
 17. Zheng C, Zeng D, Chen J, Liu S, et al. Effectiveness of extracorporeal shock wave therapy in patients with tennis elbow: A meta-analysis of randomized controlled trials. *Medicine (Baltimore).* 2020;99(30):e21189.
 18. Yoon SY, Kim YW, Shin IS, Moon HI, et al. Does the type of extracorporeal shock therapy influence treatment effectiveness in lateral epicondylitis? A systematic review and meta-analysis. *Clin Orthop Relat Res.* 2020;478(10):2324-2339.
 19. Yao G, Chen J, Duan Y, Chen X. Efficacy of extracorporeal shock wave therapy for lateral epicondylitis: A systematic review and meta-analysis. *Biomed Res Int.* 2020;2020:2064781.
 20. Kvalvaag E, Roe C, Engebretsen KB, Soberg HL, et al. One year results of a randomized controlled trial on radial extracorporeal shock wave treatment, with predictors of pain, disability and return to work in patients with subacromial pain syndrome. *Eur J Phys Rehabil Med.* 2018;54(3):341-350.
 21. Lynen N, De Vroey T, Spiegel I, Van Ongeval F, et al. Comparison of peritendinous hyaluronan injections versus extracorporeal shock wave therapy in the treatment of painful achilles' tendinopathy: A randomized clinical efficacy and safety study. *Arch Phys Med Rehabil.* 2017;98(1):64-71.
 22. Thijs KM, Zwerver J, Backx FJ, Steeneken V, et al. Effectiveness of shockwave treatment combined with eccentric training for patellar tendinopathy: A double-blinded randomized study. *Clin J Sport Med.* 2017;27(2):89-96.

23. Newman P, Waddington G, Adams R. Shockwave treatment for medial tibial stress syndrome: A randomized double blind sham-controlled pilot trial. *J Sci Med Sport*. 2017;20(3):220-224.
24. Zhang L, Fu XB, Chen S, Zhao ZB, et al. Efficacy and safety of extracorporeal shock wave therapy for acute and chronic soft tissue wounds: A systematic review and meta-analysis. *Int Wound J*. 2018;15(4):590-599.
25. Russe E, Wechselberger G, Schwaiger K, Heinrich K, et al. Effects of preoperative extracorporeal shockwave therapy on scar formation-a pilot study on 24 subjects undergoing abdominoplasty surgery. *Lasers Surg Med*. 2020;52(2):159-165.
26. Omar MT, Gwada RF, Shaheen AA, Saggini R. Extracorporeal shockwave therapy for the treatment of chronic wound of lower extremity: Current perspective and systematic review. *Int Wound J*. 2017;14(6):898-908.
27. Korakakis V, Whiteley R, Tzavara A, Malliaropoulos N. The effectiveness of extracorporeal shockwave therapy in common lower limb conditions: A systematic review including quantification of patient-rated pain reduction. *Br J Sports Med*. 2018;52(6):387-407.
28. Cooper B, Bachoo P. Extracorporeal shock wave therapy for the healing and management of venous leg ulcers. *Cochrane Database Syst Rev*. 2018;6(6):CD011842.
29. Aguilera-Sáez J, Muñoz P, Serracanta J, Monte A, et al. Extracorporeal shock wave therapy role in the treatment of burn patients. A systematic literature review. 2019:S0305-4179(19)30211-6.
30. Karanasios S, Tsamasiotis GK, Michopoulos K, Sakellari V, Gioftsos G. Clinical effectiveness of shockwave therapy in lateral elbow tendinopathy: Systematic review and meta-analysis. *Clin Rehabil*. 2021;35(10):1383-1398.
31. Pinitkwamdee S, Laohajaroensombat S, Orapin J, Woratanarat P. Effectiveness of extracorporeal shockwave therapy in the treatment of chronic insertional achilles tendinopathy. *Foot Ankle Int*. 2020;41(4):403-410.