

References

G-43

1. Gao J, Zhang M, Zhou L, et al. Correlation between fractional exhaled nitric oxide and sputum eosinophilia in exacerbations of COPD. *International Journal of Chronic Obstructive Pulmonary Disease*.
2. Lu M, Wu B, Che D, Qiao R, Gu H. FeNO and Asthma Treatment in Children: A Systematic Review and Meta-Analysis. Wang. Y, ed. *Medicine*.
3. Matsunaga K, Hirano T, Oka A, et al. Persistently high exhaled nitric oxide and loss of lung function in controlled asthma. *Allergol Int*. Jul 2016; 65(3):266-271.
4. Blake TL, Chang AB, Chatfield MD, et al. Does Ethnicity Influence Fractional Exhaled Nitric Oxide in Healthy Individuals? A Systematic Review. *Chest*. Jul 2017;
5. Silkoff P, Laviolette M, Loza M, et al. Longitudinal stability of asthma characteristics and biomarkers from the Airways Disease Endotyping for Personalized Therapeutics (ADEPT) study. *Respiratory Research*. . Available from: MEDLINE Complete, Ipswich, MA.
6. Hayes, Inc. Hayes Medical Technology Directory Report. *Title of Report*. Lansdale, PA: Hayes, Inc.; Month 10, Year 2018.
7. Mostafavi-Pour-Manshadi S-M-Y, Naderi N, Barrechehuren M, Dehghan A, Bourbeau J. Investigating Fractional Exhaled Nitric Oxide in Chronic Obstructive Pulmonary Disease (COPD) and Asthma-COPD Overlap (ACO): A Scoping Review. *COPD: Journal of Chronic Obstructive Pulmonary Disease*. 2018;15(4):377-391.