

References

L-194

1. Sanghi V and Thota PN. Barrett's esophagus: Novel strategies for screening and surveillance. *Ther Adv Chronic Dis.* 2019;10:1-14.
2. Kaul V, Gross S, Corbett FS,. Clinical utility of wide-area transepithelial sampling with three-dimensional computer-assisted analysis (WATS3D) in identifying Barrett's esophagus and associated neoplasia. *Dis Esophagus.* 2020;33(12):doaa069.
3. Docimo S Jr, Al-Mansour M, Tsuda S. SAGES TAVAC safety and efficacy analysis WATS3D (CDx Diagnostics, Suffern, NY). *Surg Endosc.* 2020;34(9):3743-3747.
4. Singer ME, Smith MS. Wide area transepithelial sampling with computer-assisted analysis (WATS3D) is cost-effective in Barrett's esophagus screening. *Dig Dis Sci.* 2021;66(5):1572-1579.
5. Agha YH, Taleb A, Srinivasan S, et al. Screening for Barrett's esophagus in patients with cirrhosis using WATS3D. *Kans J Med.* 2021;14:206-208.
6. Raphael KL, Stewart M, Sejpal DV, et al. Adjunctive yield of wide-area transepithelial sampling for dysplasia detection after advanced imaging and random biopsies in Barrett's esophagus. *Clin Transl Gastroenterol.* 2019;10(12):e00107.
7. Smith MS, Ikonomi E, Bhuta R, et al; US Collaborative WATS Study Group. Wide-area transepithelial sampling with computer-assisted 3-dimensional analysis (WATS) markedly improves detection of esophageal dysplasia and Barrett's esophagus: Analysis from a prospective multicenter community-based study. *Dis Esophagus.* 2019;32(3):doy099.
8. Agha YH, Srinivasan S, Hyder J, et al. WATS3D versus forceps biopsy in screening for Barrett's esophagus: Experience in community endoscopy centers. *Ann Gastroenterol.* 2021;34(2):164-168.
9. Odze RD, Goldblum J, Kaul V. Role of wide-area transepithelial sampling with 3D computer-assisted analysis in the diagnosis and management of Barrett's Esophagus. *Clin Transl Gastroenterol.* 2021;12(12):e00422.
10. Codipilly DC, Krishna Chandar A, Wang KK, et al. Wide-area transepithelial sampling for dysplasia detection in Barrett's esophagus: A systematic review and meta-analysis. *Gastrointest Endosc.* 2022;95(1):51-59.e7.
11. Shaheen NJ, Smith MS, Odze RD. Progression of Barrett's esophagus, crypt dysplasia, and low-grade dysplasia diagnosed by wide-area transepithelial sampling with 3-dimensional computer-assisted analysis: A retrospective analysis. *Gastrointest Endosc.* 2022;95(3):410-418.e1.

12. Agha YH, Srinivasan S, Hyder J, et al. WATS 3D versus forceps biopsy in screening for Barrett's esophagus: Experience in community endoscopy centers. *Ann Gastroenterol*. 2021;34(2):164-168.
13. CDx Diagnostics. WATS3D. 2021; <https://www.cdxdiagnostics.com/wats3d>. Accessed July 2, 2021.
14. ASGE STANDARDS OF PRACTICE COMMITTEE, Qumseya B, Sultan S, et al. ASGE guideline on screening and surveillance of Barrett's esophagus. *Gastrointest Endosc*. 2019;90(3):335-359.e2.
15. Souza RF, Spechler SJ. Advances in biomarkers for risk stratification in Barrett's esophagus. *Gastrointest Endosc Clin N Am*. 2021;31(1):105-115.