## References

## **M-50**

- 1. Frontera A, Carpenter A, Duncan E, et al. Demographic and clinical characteristics to predict paroxysmal atrial fibrillation: insights from an implantable loop recorder population. *Pacing Clin Electrophysiol*. 2015; 38(10):1217-1222.
- 2. Ergul Y, Tanidir I, Ozyilmaz I, Akdeniz C, Tuzcu V. Evaluation rhythm problems in unexplained syncope etiology with implantable loop recorder. *Pediatr Int*. 2015; 57(3):359-366. Accessed June 9, 2017.
- 3. Afzal, M.R., Gunda, S., Waheed, S., et al. Role of outpatient cardiac rhythm monitoring in cryptogenic stroke: a systematic review and metaanalysis. *Pacing Clin Electrophysiol*. 2015;38(10):1236-1245. Accessed June 14, 2017.
- 4. Poli S, Diedler J, Zuern C, et al. Insertable cardiac monitors after cryptogenic stroke--a risk factor based approach to enhance the detection rate for paroxysmal atrial fibrillation. *Euro J Neurol*. 2016; 23(2):375-381. Accessed June 14, 2017.
- 5. Podrid P.J. Ambulatory ECG Monitoring. In: UpToDate, Post TW (Ed), UpToDate, Waltham, MA. Accessed on June 6, 2016
- 6. Brachmann J., Morillo C. A., Sanna T., et al. Uncovering atrial fibrillation beyond short-term monitoring in cryptogenic stroke patients three-year results from the cryptogenic stroke and underlying atrial fibrillation trial. *Circ Arrhyth Electrophysiol*. 2016; 9.
- 7. Galli A, Ambrosini F, Lombardi F. Holter monitoring and loop recorders: from research to clinical practice. *Arrhyth Electrophysiol Rev.* 2016; 5(2):136-143.
- Bolourchi M, Batra A. Diagnostic yield of patch ambulatory electrocardiogram monitoring in children (from a national registry). *Am J Cardiol*. 2015; 115(5):630-634. Accessed June 19, 2017.
- 9. Experts release guidelines for evaluating, managing syncope. *News*. 2017. Accessed June 6, 2017.

- 10. Mobile cardiac outpatient telemetry for detecting arrhythmias. ECRI Institute. 2015. Accessed February 24, 2017.
- 11. Solomon M, Yang J, Sung S, et al. Incidence and timing of potentially highrisk arrhythmias detected through long term continuous ambulatory electrocardiographic monitoring. *BMC Cardiovasc Dis*. 2016; 16:35.
- 12. Sangha, R.S., Bernstein, R. Advances in the detection and monitoring of atrial fibrillation for patients with cryptogenic ischemic stroke. *Curr Atheroscler Rep* (2015) 17: 71.
- 13. January CT, Wann LS, Alpert JS, et al. 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: A report of the American college of cardiology/American heart association task force on practice guidelines and the heart rhythm Society. *J Am Coll Cardiol*. 201; 64(21):e1-e76.
- 14. Hayes, Inc. Hayes Health Technology Brief. Implantable cardiac loop recorders for detection of atrial fibrillation following cryptogenic stroke. Lansdale, PA: Hayes, Inc.; Published on 06/15/2017. Reviewed on 06/21/2018. Accessed on 07/16/2019.
- 15. Hayes, Inc. Hayes Health Technology Brief. Zio Patch (iRhythm Technologies, Inc.) long-term ambulatory cardiac rhythm monitoring. Lansdale, PA: Hayes, Inc.; Published on 02/07/2019. Accessed on 07/16/2019.
- 16. Sheikh AB, Felzer JR, Munir AB, Morin DP, Lavie CJ. Evaluating the benefits of home-based management of atrial fibrillation: Current perspectives. Pragmat Obs Res. 2016; 7:41 53.