

References

M-70

1. Centers for Medicare and Medicaid Services (CMS). National Coverage Determination (NCD). 20.19: Ambulatory blood pressure monitoring. [CMS Web site]. 07/01/03.
2. Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: Executive summary: A report of the American College of Cardiology/American Heart Association Task Force on clinical practice guidelines. *J Am Coll Cardiol*. 2018;71(19):2199-2269.
3. Kupferman J, Batisky D, Samuels J, et al. Ambulatory blood pressure monitoring and neurocognitive function in children with primary hypertension. *Pediatr Nephrol* 2018;33:1765–1771.
4. National Institute for Health and Care Excellence. Hypertension in adults: Diagnosis and management [NG136].2019; <https://www.nice.org.uk/guidance/ng1>
5. Muntner P, Shimbo D, Carey RM, et al. Measurement of Blood Pressure in Humans: A scientific statement from the American Heart Association. 2019;73(5):e35- e66.
6. Yang WY, Melgarejo JD, Thijs L, et al. Association of office and ambulatory blood pressure with mortality and cardiovascular outcomes. 2019;322(5):409-420.
7. Hayes, Inc. Hayes Health Technology. *Zio Patch (iRhythm Technologies Inc.) Long-term ambulatory cardiac rhythm monitoring*. Lansdale, PA. Hayes, Inc.; 02/07/2019.
8. Aung K, Htay T. Relationship between outpatient clinic and ambulatory blood pressure measurements and mortality. *Curr Cardiol Rep*. 2019;21(5):28.