

## References

### Z-64

1. Hilmisson H, Berman S, Magnúsdóttir S. Sleep apnea diagnosis in children using software-generated apnea-hypopnea index (AHI) derived from data recorded with a single photoplethysmogram sensor (PPG) : Results from the Childhood Adenotonsillectomy Study (CHAT) based on cardiopulmonary coupling analysis [published online ahead of print, 2020 Mar 28]. *Sleep Breath*. 2020;10.1007/s11325-020-02049-6.
2. Garde A, Hoppenbrouwer X, Dehkordi P, et al. Pediatric pulse oximetry-based OSA screening at different thresholds of the apnea-hypopnea index with an expression of uncertainty for inconclusive classifications. *Sleep Med*. 2019;60:45-52.
3. Mitchell RB, Archer SM, Ishman SL, et al. Clinical Practice Guideline: Tonsillectomy in Children (Update)-Executive Summary. *Otolaryngol Head Neck Surg*. 2019;160(2):187-205.
4. Rosenberg R, Hirshkowitz M, Rapoport DM, Kryger M. The role of home sleep testing for evaluation of patients with excessive daytime sleepiness: Focus on obstructive sleep apnea and narcolepsy. *Sleep Med*. 2019;56:80-89.
5. Lin CH, Chen CH, Hong SY, Chou IC, Liang SJ, Hang LW. Polysomnography is an important method for diagnosing pediatric sleep problems: Experience of one children's hospital. *Children (Basel)*. 2021;8(11):991.
6. Anitua E, Mayoral P, Almeida GZ, Durán-Cantolla J, Alkhraisat MH. A multicenter prospective study on the use of a mandibular advancement device in the treatment of obstructive sleep apnea. *Dentistry J*. 2023; 11(11):247.
7. Nosetti L, Zaffanelli M, Katz E et al, Twenty-year follow-up of children with obstructive sleep apnea. 2022;18(6):1573-1581.
8. Yu P, Radcliff J, Taylor H, et al. Neurobehavioral morbidity of pediatric mild sleep-disordered breathing and obstructive sleep apnea. 2022;45(5):zsac035