

References

S-31

1. Spaans A, van Minnen L, Kon M, et al. Conservative treatment of thumb base osteoarthritis: a systematic review. *J Hand Surg Am.* 2015;40(1):16-21.e1-6.
2. Wu T, Dong Y, Song H, et al. Ultrasound-guided versus landmark in knee arthrocentesis: A systematic review. *Semin. Arthritis Rheum.* 2016;627-632.
3. Rice DA, McNair PJ, Lewis GN, et al. The effects of joint aspiration and intra-articular corticosteroid injection on flexion reflex excitability, quadriceps strength and pain in individuals with knee synovitis: a prospective observational study. *Arthritis Res Ther.* 2015;17:191.
4. Boelch SP, Roth M, Arnholdt J, et al. Synovial fluid aspiration should not be routinely performed during the two-stage exchange of the knee. *Biomed Res Int.* 2018:1-8.
5. Gurung T, Singh RK, Mohammad S, et al. Efficacy of arthrocentesis versus arthrocentesis with sodium hyaluronic acid in temporomandibular joint osteoarthritis: A comparison. *Natl J Maxillofac Surg.* 2017;8(1):41.
6. Yaqub S, Sibbitt Jr WL, Band PA, et al. Can diagnostic and therapeutic arthrocentesis be successfully performed in the flexed knee? *J Clin Rheumatol.* 2018;24(6):295-301.
7. Bhavsar TB, Sibbitt WL, Band PA, et al. Improvement in diagnostic and therapeutic arthrocentesis via constant compression. *Clin Rheumatol.* 2018;37(8):2251-9.