## References

## **O-13**

- 1. Ellwood J, Draper-Rodi J, Carnes D. The effectiveness and safety of conservative interventions for positional plagiocephaly and congenital muscular torticollis: A synthesis of systematic reviews and guidance. *Chiropr Man Therap.* 2020;28(1):1-11.
- 2. Tamber MS, Nikas D, Beier A, et al. The role of cranial molding orthosis (helmet) therapy. 2016; https://www.cns.org/guidelines/browse-guidelines-detail/5-role-of-cranial-molding-orthosis-helmet-therapy. Accessed January 12, 2022.
- 3. National Institute of Neurological Disorders and Stroke (NINDS). Craniosynostosis Information Page. 2017; https://www.ninds.nih.gov/Disorders/All-Disorders/Craniosynostosis-Information-Page. Accessed January 28, 2021.
- 4. Graham T, Adams-Huet B, Gilbert N, Witthoff K, Gregory T, Walsh M. Effects of initial age and severity on cranial remolding orthotic treatment for infants with deformational plagiocephaly. *J Clin Med.* 2019;8(8):1097.
- 5. Kunz, F., Schweitzer, T., Große, S., et.al. Head orthosis therapy in positional plagiocephaly: Longitudinal 3D-investigation of long-term outcomes, compared with untreated infants and with a control group. *Eur J Orthod.* 2019;41(1):29-37.
- 6. Cevik S, Isik S, Oskilic A, The role of age on helmet therapy in deformational plagiocephaly and asymmetric brachycephaly. *Childs Nerv Syst.* 2019;36:803-810.
- 7. Marino s, Ruggieri M, Marino L, Falsaperla R. Suture ultrasound: Useful diagnostic screening for posterior plagiocephaly. *Childs Nerv Syst*. 2021;37:3715-3720.