

1. Boswell MV, Manchikanti L, Kaye AD, et al. A best-evidence systematic appraisal of the diagnostic accuracy and utility of facet (zygapophysial) joint injections in chronic spinal pain. *Pain Physician*. Jul-Aug 2015;18(4):E497-533. PMID 26218947.
2. Falco FJ, Datta S, Manchikanti L, et al. An updated review of the diagnostic utility of cervical facet joint injections. *Pain Physician*. Nov-Dec 2012;15(6):E807-838. PMID 23159977.
3. Falco FJ, Manchikanti L, Datta S, et al. Systematic review of the therapeutic effectiveness of cervical facet joint interventions: an update. *Pain Physician*. Nov-Dec 2012;15(6):E839-868. PMID 23159978.
4. Falco FJ, Manchikanti L, Datta S, et al. An update of the systematic assessment of the diagnostic accuracy of lumbar facet joint nerve blocks. *Pain Physician*. Nov-Dec 2012;15(6):E869-907. PMID 23159979.
5. Falco FJ, Manchikanti L, Datta S, et al. An update of the effectiveness of therapeutic lumbar facet joint interventions. *Pain Physician*. Nov-Dec 2012;15(6):E909-953. PMID 23159980.
6. Cohen SP, Strassels SA, Kurihara C, et al. Randomized study assessing the accuracy of cervical facet joint nerve (medial branch) blocks using different injectate volumes. *Anesthesiology*. Jan 2010;112(1):144-152. PMID 19996954.
7. Cohen SP, Stojanovic MP, Crooks M, et al. Lumbar zygapophysial (facet) joint radiofrequency denervation success as a function of pain relief during diagnostic medial branch blocks: a multicenter analysis. *Spine J*. May- Jun 2008;8(3):498-504. PMID 17662665.
8. Pampati S, Cash KA, Manchikanti L. Accuracy of diagnostic lumbar facet joint nerve blocks: a 2-year follow-up of 152 patients diagnosed with controlled diagnostic blocks. *Pain Physician*. Sep-Oct 2009;12(5):855-866. PMID 19787011.
9. Manchikanti L, Pampati S, Cash KA. Making sense of the accuracy of diagnostic lumbar facet joint nerve blocks: an assessment of the implications of 50% relief, 80% relief, single block, or controlled diagnostic blocks. *Pain Physician*. Mar-Apr 2010;13(2):133-143. PMID 20309379.
10. Manchikanti L, Kaye AD, Boswell MV, et al. A systematic review and best evidence synthesis of the effectiveness of therapeutic facet joint interventions in managing chronic spinal pain. *Pain Physician*. Jul-Aug 2015;18(4):E535-582. PMID 26218948.
11. Chou R, Atlas SJ, Stanos SP, et al. Nonsurgical interventional therapies for low back pain: a review of the evidence for an American Pain Society clinical practice guideline. *Spine (Phila Pa 1976)*. May 1 2009;34(10):1078-1093. PMID 19363456.

12. Civelek E, Cansever T, Kabatas S, et al. Comparison of effectiveness of facet joint injection and radiofrequency denervation in chronic low back pain. *Turk Neurosurg.* Mar 2012;22(2):200-206. PMID 22437295.
13. Lakemeier S, Lind M, Schultz W, et al. A comparison of intraarticular lumbar facet joint steroid injections and lumbar facet joint radiofrequency denervation in the treatment of low back pain: a randomized, controlled, double-blind trial. *Anesth Analg.* Jul 2013;117(1):228-235. PMID 23632051.
14. Nath S, Nath CA, Pettersson K. Percutaneous lumbar zygapophysial (facet) joint neurotomy using radiofrequency current, in the management of chronic low back pain: a randomized double-blind trial. *Spine (Phila Pa 1976).* May 20 2008;33(12):1291-1297; discussion 1298. PMID 18496338.
15. Haspeslagh SR, Van Suijlekom HA, Lame IE, et al. Randomised controlled trial of cervical radiofrequency lesions as a treatment for cervicogenic headache [ISRCTN07444684]. *BMC Anesthesiol.* Feb 16 2006;6:1. PMID 16483374.
16. van Wijk RM, Geurts JW, Wynne HJ, et al. Radiofrequency denervation of lumbar facet joints in the treatment of chronic low back pain: a randomized, double-blind, sham lesion-controlled trial. *Clin J Pain.* Jul-Aug 2005;21(4):335-344. PMID 15951652.
17. Lord SM, Barnsley L, Wallis BJ, et al. Percutaneous radio-frequency neurotomy for chronic cervical zygapophyseal-joint pain. *N Engl J Med.* Dec 5 1996;335(23):1721-1726. PMID 8929263.
18. Husted DS, Orton D, Schofferman J, et al. Effectiveness of repeated radiofrequency neurotomy for cervical facet joint pain. *J Spinal Disord Tech.* Aug 2008;21(6):406-408. PMID 18679094.
19. Schofferman J, Kine G. Effectiveness of repeated radiofrequency neurotomy for lumbar facet pain. *Spine (Phila Pa 1976).* Nov 1 2004;29(21):2471-2473. PMID 15507813.
20. Smuck M, Crisostomo RA, Trivedi K, et al. Success of initial and repeated medial branch neurotomy for zygapophysial joint pain: a systematic review. *PM R.* Sep 2012;4(9):686-692. PMID 22980421.
21. Rambaransingh B, Stanford G, Burnham R. The effect of repeated zygapophysial joint radiofrequency neurotomy on pain, disability, and improvement duration. *Pain Med.* Sep 2010;11(9):1343-1347. PMID 20667024.
22. Manchikanti L, Singh V, Falco FJ, et al. Comparative outcomes of a 2-year follow-up of cervical medial branch blocks in management of chronic neck pain: a randomized, double-blind controlled trial. *Pain Physician.* Sep-Oct 2010;13(5):437-450. PMID 20859313.
23. Manchikanti L, Singh V, Falco FJ, et al. Evaluation of lumbar facet joint nerve blocks in managing chronic low back pain: a randomized, double-blind,

- controlled trial with a 2-year follow-up. *Int J Med Sci.* May 28 2010;7(3):124-135. PMID 20567613.
24. Manchikanti L, Singh V, Falco FJ, et al. Comparative effectiveness of a one-year follow-up of thoracic medial branch blocks in management of chronic thoracic pain: a randomized, double-blind active controlled trial. *Pain Physician.* Nov-Dec 2010;13(6):535-548. PMID 21102966.
  25. Manchikanti L, Singh V, Falco FJ, et al. The role of thoracic medial branch blocks in managing chronic mid and upper back pain: a randomized, double-blind, active-control trial with a 2-year followup. *Anesthesiol Res Pract.* 2012;2012:585806. PMID 22851967.
  26. Hashemi M, Hashemian M, Mohajerani SA, et al. Effect of pulsed radiofrequency in treatment of facet-joint origin back pain in patients with degenerative spondylolisthesis. *Eur Spine J.* Sep 2014;23(9):1927-1932. PMID 24997616.
  27. Kroll HR, Kim D, Danic MJ, et al. A randomized, double-blind, prospective study comparing the efficacy of continuous versus pulsed radiofrequency in the treatment of lumbar facet syndrome. *J Clin Anesth.* Nov 2008;20(7):534-537. PMID 19041042.
  28. Van Zundert J, Patijn J, Kessels A, et al. Pulsed radiofrequency adjacent to the cervical dorsal root ganglion in chronic cervical radicular pain: a double blind sham controlled randomized clinical trial. *Pain.* Jan 2007;127(1- 2):173-182. PMID 17055165.
  29. Tekin I, Mirzai H, Ok G, et al. A comparison of conventional and pulsed radiofrequency denervation in the treatment of chronic facet joint pain. *Clin J Pain.* Jul-Aug 2007;23(6):524-529. PMID 17575493.
  30. Iwatsuki K, Yoshimine T, Awazu K. Alternative denervation using laser irradiation in lumbar facet syndrome. *Lasers Surg Med.* Mar 2007;39(3):225-229. PMID 17345622.
  31. Joo YC, Park JY, Kim KH. Comparison of alcohol ablation with repeated thermal radiofrequency ablation in medial branch neurotomy for the treatment of recurrent thoracolumbar facet joint pain. *J Anesth.* Jun 2013;27(3):390-395. PMID 23192698.
  32. Haufe SM, Mork AR. Endoscopic facet debridement for the treatment of facet arthritic pain--a novel new technique. *Int J Med Sci.* May 25 2010;7(3):120-123. PMID 20567612.
  33. Watters WC, 3rd, Resnick DK, Eck JC, et al. Guideline update for the performance of fusion procedures for degenerative disease of the lumbar spine. Part 13: injection therapies, low-back pain, and lumbar fusion. *J Neurosurg Spine.* Jul 2014;21(1):79-90. PMID 24980590.
  34. Manchikanti L, Abdi S, Atluri S, et al. An update of comprehensive evidence-based guidelines for interventional techniques in chronic spinal pain. Part II:

guidance and recommendations. *Pain Physician*. Apr 2013;16(2 Suppl):S49-283. PMID 23615883.

35. American Society of Anesthesiologists Task Force on Chronic Pain Management, American Society of Regional Anesthesia and Pain Medicine. Practice guidelines for chronic pain management: an updated report by the American Society of Anesthesiologists Task Force on Chronic Pain Management and the American Society of Regional Anesthesia and Pain Medicine. *Anesthesiology*. Apr 2010;112(4):810-833. PMID 20124882.
36. National Institute for Health and Clinical Excellence (NICE). NICE guideline [NG59]: Low back pain and sciatica in over 16s: assessment and management. 2016; <https://www.nice.org.uk/guidance/NG59>. Accessed September 20, 2019.
37. California Technology Assessment Forum (CTAF). Percutaneous radiofrequency neurotomy for treatment of chronic pain from the upper cervical (C2-3) spine. A Technology Assessment. 2007; [http://icer-review.org/wp-content/uploads/2016/01/742\\_file\\_Neurotomy\\_Web.pdf](http://icer-review.org/wp-content/uploads/2016/01/742_file_Neurotomy_Web.pdf). Accessed September 20, 2019.