

References

Z-75

1. Vecchioli-Scaldazza C, Morosetti C, Giampieretti R, et al. Percutaneous tibial nerve stimulation versus electrical stimulation with pelvic floor muscle training for overactive bladder syndrome in women: results of a randomized controlled study. *Int Braz J Urol.* 2017; 43(1): 121-126.
2. Stewart F, Gameiro LF, El Dib R, et al. Electrical stimulation with non-implanted electrodes for overactive bladder in adults. *Cochrane Database Syst Rev.* 2016;(12).
3. Simillis C, Lal N, Qiu S, et al. Sacral nerve stimulation versus percutaneous tibial nerve stimulation for faecal incontinence: a systematic review and meta-analysis. *International J Colorectal Dis.* 2018 ;33(5):645-8.
4. Sanagapalli S, Neilan L, Lo JYT, et al. Efficacy of percutaneous posterior tibial nerve stimulation for the management of fecal incontinence in multiple sclerosis: a pilot study. *Neuromodulation.* 2018;21(7):682-7.
5. Hayes, Inc. Comparative Effectiveness Review. *Comparative Effectiveness Review of Percutaneous Tibial Nerve Stimulation for the Treatment of Symptomatic Non-Neurogenic Overactive Bladder.* Lansdale, PA: Hayes, Inc.; February 4, 2020.
6. Hayes, Inc. Medical Technology Directory. *Percutaneous Tibial Nerve Stimulation for the Treatment of Symptomatic Neurogenic Lower Urinary Tract Dysfunction.* Lansdale, Pa: Hayes, Inc.; April 15, 2019.