

References

Z-75

1. Sarveazad A, Babahajian A, Amini N, et al. Posterior tibial nerve stimulation in fecal incontinence: A systematic review and meta-analysis. *Basic Clin Neurosci.* 2019;10(5):419-431.
2. Tan K, Wells CI, Dinning P, et al. Placebo response rates in electrical nerve stimulation trials for fecal incontinence and constipation: A systematic review and meta-analysis. *2020;23(8):1108-1116.*
4. Kavanagh A, Baverstock R, Campeau L, et al. Canadian Urological Association guideline: Diagnosis, management, and surveillance of neurogenic lower urinary tract dysfunction. *Can Urol Assoc J.* 2019;13(6):E157-E176.
5. Rahnama'i MS. Neuromodulation for functional bladder disorders in patients with multiple sclerosis. *Mult Scler.* 2020;26(11):1274-1280.
5. Lane GI, Mao-Draayer Y, Barboglio-Romo P, et al. A prospective observational cohort study of posterior tibial nerve stimulation in patients with multiple sclerosis: Design and methods. *BMC Urol.* 2020;20(1):58.
6. Tudor KI, Seth JH, Liechti MD, et al. Outcomes following percutaneous tibial nerve stimulation (PTNS) treatment for neurogenic and idiopathic overactive bladder. *Clin Auton Res.* 2020;30(1):61-67.
7. Andersen K, Kobberø H, Pedersen TB, Poulsen MH. Percutaneous tibial nerve stimulation for idiopathic and neurogenic overactive bladder dysfunction: A four-year follow-up single-centre experience. *Scand J Urol.* 2021;55(2):169-176.
8. Wang M, Jian Z, Ma Y, Jin X, Li H, Wang K. Percutaneous tibial nerve stimulation for overactive bladder syndrome: A systematic review and meta-analysis. *Int Urogynecol J.* 2020;31(12):2457-2471.
9. Xiong SC, Peng L, Hu X, Shao YX, Wu K, Li X. Effectiveness and safety of tibial nerve stimulation versus anticholinergic drugs for the treatment of overactive bladder syndrome: A meta-analysis. *Ann Palliat Med.* 2021;10(6):6287-6296.

10. Lo CW, Wu MY, Yang SS, Jaw FS, Chang SJ. Comparing the efficacy of OnabotulinumtoxinA, sacral neuromodulation, and peripheral tibial nerve stimulation as third line treatment for the management of overactive bladder symptoms in adults: Systematic review and network meta-analysis. *Toxins (Basel)*. 2020 Feb 18;12(2):128.
11. Lashin AM, Eltabey NA, Wadie BS. Percutaneous tibial nerve stimulation versus sham efficacy in the treatment of refractory overactive bladder: Outcomes following a shortened 6-week protocol, a prospective randomized controlled trial. *Int Urol Nephrol*. 2021;53(12):2459-2467.
12. Vollstedt A, Gilleran J. Update on implantable PTNS devices. *Curr Urol Rep*. 2020;21(7):28.
13. Yamashiro J, de Riese W, de Riese C. New implantable tibial nerve stimulation devices: Review of published clinical results in comparison to established neuromodulation devices. *Res Rep Urol*. 2019;11:351-357.
14. Te Dorsthorst M, Digesu A, van Kerrebroeck P, et al. Patient-tailored healthcare and tibial nerve neuromodulation in the treatment of patients with overactive bladder symptoms. *Neurourol Urodyn*. 2022; 41(2):679-684.
15. Coolen RL, Groen J, Scheepe JR, et al. Transcutaneous electrical nerve stimulation and percutaneous tibial nerve stimulation to treat idiopathic nonobstructive urinary retention: A systematic review. *Eur Urol Focus*. 2021; 7(5): 1184-1194.
16. Ho FCS, He C, Yao HH, et al. Efficacy of sacral neuromodulation and percutaneous tibial nerve stimulation in the treatment of chronic nonobstructive urinary retention: A systematic review. *Neurourol Urodyn*. 2021; 40(5):1078-1088.
17. Rogers A, Bragg S, Ferrante K, et al. Pivotal study of leadless tibial nerve stimulation with eCoin® for urgency urinary incontinence: An open-label, single arm trial. *J Urol*. 2021;206(2): 399-408.
18. Gilling P, Meffan P, Kaaki B, et al. Twelve-month durability of a fully-implanted, nickel-sized and shaped tibial nerve stimulator for the treatment of overactive bladder syndrome with urgency urinary incontinence: A single-arm, prospective study. *Urology*. 2021;157:71-78.

19. Kaaki B, English S, Gilling P, et al. Six-month outcomes of reimplantation of a coin-sized tibial nerve stimulator for the treatment of overactive bladder syndrome with urgency urinary incontinence. *Female Pelvic Med Reconstr Surg.* 2022; 28(5): 287-292.
20. Luo C, Wei D, Pang K, et al. Is percutaneous tibial nerve stimulation (PTNS) effective for fecal incontinence (FI) in adults compared with sham electrical stimulation? A meta-analysis. *Tech Coloproctol.* 2024; 28(1):37.
21. Leo CA, Thomas GP, Hodgkinson JD, et al. Randomized pilot study: Anal inserts versus percutaneous tibial nerve stimulation in patients with fecal incontinence. *Dis Colon Rectum.* 2021; 64(4):466-474.
22. Zyczynski HM, Richter HE, Sung VW, et al. Percutaneous tibial nerve stimulation vs sham stimulation for fecal incontinence in women: NeurOmodulaTion for accidental bowel leakage randomized clinical trial. *Am J Gastroenterol.* 2022;117(4):654-667.
23. Cameron AP, Chung DE, Dielubanza EJ, et al. The AUA/SUFU guideline on the diagnosis and treatment of idiopathic overactive bladder. *J Urol.* 2024;212(1):11-20.
24. Tahmasbi F, Salehi-Pourmehr H, Naseri A, et al. Effects of posterior tibial nerve stimulation (PTNS) on lower urinary tract dysfunction: An umbrella review. *Neurourol Urodyn.* 2024;43(2):494-515.
25. Kendall HJ, Schrijvers J, Heesakkers JPFA. Current position of neuromodulation for bladder pain syndrome/interstitial cystitis. *Curr Opin Urol.* 2024;34(2):64-68.
26. Abdalla AS, Berquist SW, Mohammad A, Dobberfuhl AD. Our 11-year experience with percutaneous tibial nerve stimulation in men and women for the treatment of interstitial cystitis/bladder pain syndrome. *Urology.* 2023;182:73-78.
27. Huang J, Fan Y, Zhao K, et al. Comparative efficacy of neuromodulation technologies for overactive bladder in adults: A Network meta-analysis of randomized controlled trials. *Neuromodulation.* 2023;26(8):1535-1548.