

References

S-159

1. National Institute for Health and Care Excellence (NICE). Percutaneous coblation of the intervertebral disc for low back pain and sciatica. 2016. Accessed January 16, 2018.
2. Cincu R, Lorente F de A, Gomez J, Eiras J, Agrawal A. One decade follow up after nucleoplasty in the management of degenerative disc disease causing low back pain and radiculopathy. *Asian Journal of Neurosurgery*. 2015;10(1):21-25.
3. Ren D-J, Liu X-M, Du S-Y, Sun T-S, Zhang Z-C, Li F. Percutaneous Nucleoplasty Using Coblation Technique for the Treatment of Chronic Nonspecific Low Back Pain: 5-year Follow-up Results. *Chinese Medical Journal*. 2015;128(14):1893-1897.
4. Eichen PM, Achilles N, Kirchner R, et al. Nucleoplasty, a minimally invasive procedure for disc decompression: a systematic review and meta-analysis of published clinical studies. *Pain Physician [serial online]*. 2014;17(2):E149-E173.
5. Yang B, Xie J, Yin B, Wang L, Fang S, et al. Treatment of cervical disc herniation through percutaneous minimally invasive techniques. *Eur Spine J*. 2014;23(2):382-388.
6. Desai MJ, Kapural L, Petersohn JD, Vallejo R, Menzies R, et al. Twelve-month follow-up of a randomized clinical trial comparing intradiscal biacuplasty to conventional medical management for discogenic lumbar back pain. *Pain Med*. 2017;18(4):751-763.