

References

Z-24

1. Levenhagen K, Davies C, Perdomo M, et al. Diagnosis of upper quadrant lymphedema secondary to cancer: Clinical practice guideline from the oncology section of the American Physical Therapy Association. *Rehabil Oncol*. 2017;35(3):E1–E18.
2. National Comprehensive Cancer Network (NCCN) Clinical Practice Guidelines in Oncology. Breast Cancer. Version 2.2022.
3. Hayes, Inc. Hayes Health Technology Assessment. Bioelectrical Impedance (Bioimpedance) Analysis for Assessment of Lymphedema. Lansdale, PA: Hayes, Inc.; 08/06/2020.
4. Sickle Cell Society. Iron Monitoring and Management in Sickle Cell Disease.
5. Centers for Disease Control and Prevention. Chelation Therapy.
6. Skróder H, Kippler M, Nermell B, et al. Major limitations in using element concentrations in hair as biomarkers of exposure to toxic and essential trace elements in children. *Environ Health Perspect*. 2017;125(6):1-9.
7. Katz SA. On the use of hair analysis for assessing arsenic intoxication. *Int J Environ Res Public Health*. 2019;16(6).
8. Da Dalt L, Parri N, Amigoni A, Nocerino A, Selmin F, et al. Italian guidelines on the assessment and management of pediatric head injury in the emergency department. *Ital J Pediatr*. 2018;44(71);1-41.
9. Panthi S, Khatri K, Kharel K, et al. Outcome of percutaneous release of tennis elbow: A nonrandomized controlled trial Study. 2017;9(1):e952.
10. Gaspar MP, Motto MA, Lewis S, Jacoby SM, Culp RW, et al. Platelet-rich plasma injection with percutaneous needling for recalcitrant lateral epicondylitis: Comparison of tenotomy and fenestration techniques. *Orthop J Sports Med*. 2017;5(12).
11. Chen C, Ma YH, Zhang YT, Zhang F, Zhou N, Wang X, et al. Effect of dendritic cell-based immunotherapy on hepatocellular carcinoma: A systematic review and meta-analysis. *Cytotherapy*. 2018;20(8):975-989.
12. Dafni U, Michielin O, Lluesma SM, Tsourti Z, Polydoropoulou V, Karlis D, et al. Efficacy of adoptive therapy with tumor-infiltrating lymphocytes and recombinant interleukin-2 in advanced cutaneous melanoma: A systematic review and meta-analysis. *Ann Oncol*. 2019;30(12):1902-1913.
13. Khammari A, Nguyen JM, Leccia MT, Guillot B, Saiagh S, Pandolfino MC, et al. Tumor infiltrating lymphocytes as adjuvant treatment in stage III melanoma patients with only one invaded lymph node after complete resection: Results from a multicentre, randomized clinical phase III trial. *Cancer Immunol Immunother*. 2020;69(8):1663-1672.
14. Cheung XC, Fahey T, Rogers AC, Pemberton JH, Kavanagh DO. Surgical management of idiopathic perianal fistulas: A systematic review and meta-analysis. *Dig Surg*. 2021;38(2):104-119.
15. Jayne DG, Scholefield J, Tolan D, Gray R, Senapati A, Hulme CT, et al.; FIAT Trial Collaborative Group. A multicenter randomized controlled trial comparing safety, efficacy, and cost-effectiveness of the surgis anal fistula plug versus surgeon's preference for transsphincteric fistula-in-ano: The FIAT Trial. *Ann Surg*. 2021;273(3):433-441.
16. Ridner SH, Dietrich MS, Cowher MS, Taback B, McLaughlin S, Ajkay N, et al. A randomized trial evaluating bioimpedance spectroscopy versus tape measurement for the prevention of

lymphedema following treatment for breast cancer: Interim analysis. *Ann Surg Oncol*. 2019;26(10):3250-3259.

17. American Diabetes Association. 9. Pharmacologic approaches to glycemic treatment: Standards of medical care in diabetes-2021. *Diabetes Care*. 2021;44(Suppl 1):S111-S124.
18. American Diabetes Association Professional Practice Committee; American Diabetes Association Professional Practice Committee.; Draznin B, Aroda VR, Bakris G, Benson G, Brown FM, Freeman R, et al. 9. Pharmacologic approaches to glycemic treatment: Standards of medical care in diabetes-2022. *Diabetes Care*. 2022;45(Suppl 1):S125-S143.
19. Bhattacharyya N, Gubbels SP, Schwartz SR, Edlow JA, El-Kashlan H, Fife T, Holmberg JM, et al. Clinical practice guideline: Benign paroxysmal positional vertigo (update). *Otolaryngol Head Neck Surg*. 2017;156(3_suppl):S1-S47.
20. Hamlin K, Munro C, Barker SL, McKenna S, Kumar K. Open release versus radiofrequency microtenotomy in the treatment of lateral epicondylitis: A prospective randomized controlled trial. *Shoulder Elbow*. 2018;10(1):45-51.
21. Wang W, Rikhranj IS, Chou ACC, Chong HC, Koo KOT. Endoscopic plantar fasciotomy vs open radiofrequency microtenotomy for recalcitrant plantar fasciitis. *Foot Ankle Int*. 2018;39(1):11-17.
22. Forstenpointner J, Ruscheweyh R, Attal N, Baron R, Bouhassira D, Enax-Krumova EK, Finnerup NB, et al. No pain, still gain (of function): The relation between sensory profiles and the presence or absence of self-reported pain in a large multicenter cohort of patients with neuropathy. 2021;162(3):718-727.
23. Azzopardi K, Gatt A, Chockalingam N, Formosa C. Hidden dangers revealed by misdiagnosed diabetic neuropathy: A comparison of simple clinical tests for the screening of vibration perception threshold at primary care level. *Prim Care Diabetes*. 2018;12(2):111-115.