References

Z-26

- 1. Nowak-Wegrzyn A, Chehade M, Groetch ME, et al. International consensus guidelines for the diagnosis and management of food protein-induced enterocolitis syndrome: Executive summary-Workgroup Report of the Adverse Reactions to Foods Committee, American Academy of Allergy, Asthma & Immunology. *J Allergy Clin Immunol.* 2017;139(4):1111-1126.
- 2. Scadding GW, Calderon MA, Shamji MH, et al. Immune Tolerance Network GRASS Study Team. Effect of 2 years of treatment with sublingual grass pollen immunotherapy on nasal response to allergen challenge at 3 years among patients with moderate to severe seasonal allergic rhinitis: The GRASS randomized clinical trial. JAMA 2017;317(6):615-625.
- 3. Slovick A, Douiri A, Muir R, et al. Intradermal grass pollen immunotherapy increases TH2 and IgE responses and worsens respiratory allergic symptoms. *J Allergy Clin Immunol.* 2017;139(6):18301839.
- 4. Anyane-Yeboa A, Wenfei Wang, Kavitt RT. The role of allergy testing in eosinophilic esophagitis. *Gastroenterology & Hepatology*. 2018;14(8):463-469.
- 5. Gill MM, Gasner S, Banken A, et. al. Improving routine prenatal penicillin allergy testing for reported penicillin allergy. *BMJ Open Qual*. 2022;11(3):e001859.
- 6. Cansever M, Oruç Ç. What plays a role in the severity of atopic dermatitis in children? *Turk J Med Sci.* 2021;51(5):2494-2501.
- 7. Meyer R, De Koker C, Dziubak R, et. al. The challenge of home allergen reintroductions using the ladder approach in children with non-IgE mediated gastrointestinal food allergy. *Front Allergy*. 2021;2:721686.