

## References

Z-75

1. Peters KM, Carrico DJ, Wooldridge LS, et al. Percutaneous Tibial Nerve Stimulation for the Long-Term Treatment of Overactive Bladder: 3-Year Results of the STEP Study. *The Journal of Urology*. 2013;189:2194-2201.
2. Gormley EA, Lightner DJ, Burgio KL. Diagnosis and treatment of overactive bladder (Non-Neurogenic) in adults:AUA/SUFU guidelines. *American Urological Association*. 2014.
3. Gaziev G, Topazio L, Iacovelli V, et al. Percutaneous tibial nerve stimulation (PTNS) efficacy in the treatment of lower urinary tract dysfunctions: a systematic review. *BMC Urol*. 2013;13:61.
4. Blue Cross and Blue Shield Association Technology Evaluation Center (TEC). Percutaneous tibial nerve stimulation for the treatment of voiding dysfunction. TEC Assessments 2013; Volume 28.
5. Zecca C, Digesu GA, Robshaw P, et al. Maintenance percutaneous posterior nerve stimulation for refractory lower urinary tract symptoms in patients with multiple sclerosis: An open label, multicenter, prospective study. *The Journal of Urology*. 2014; 191:697-702.
6. American Urological Association. AUA/SUFU Guideline: Published 2012; Amended 2014.
7. Vecchioli-Scaldazza C, Morosetti C, Berouz A, Giannubilo W, Ferrara V. Solifenacin succinate versus percutaneous tibial nerve stimulation in women with overactive bladder syndrome: results of a randomized controlled crossover study. *Gynecologic And Obstetric Investigation*. 2013;75(4):230-234
8. Yoong W, Shah P, Dadswell R, Green L. Sustained effectiveness of percutaneous tibial nerve stimulation for overactive bladder syndrome: 2-year follow-up of positive responders. *International Urogynecology Journal*. 2013;24(5):795-799.